

Division of Environmental Health and Communicable Disease Prevention								
Section: 4.0 Diseases and Conditions Updated 7/03								
Subsection: Listeriosis	Page 1 of 9							

Listeriosis Table of Contents

Listeriosis
Fact Sheet
Record of Investigation of Enteric Infection (CD-2C) rev. 6/02



Division of Environme	ntal Health and	Communicable	Disease Prevention

Section: 4.0 Diseases and Conditions	Updated 7/03
Subsection: Listeriosis	Page 2 of 9

Listeriosis

Overview^(1,2)

For a more complete description of listeriosis, refer to the following texts:

- Control of Communicable Diseases Manual (CCDM).
- Red Book, Report of the Committee on Infectious Diseases.

Case Definition⁽³⁾

Clinical description

In adults, invasive disease caused by *Listeria monocytogenes* manifests most commonly as meningitis or bacteremia; infection during pregnancy may result in fetal loss through miscarriage or stillbirth, or neonatal meningitis or bacteremia. Other manifestations can also be observed.

Laboratory criteria for diagnosis

- Isolation of *L. monocytogenes* from a normally sterile site (e.g., blood or cerebrospinal fluid [CSF] or, less commonly, joint, pleural, or pericardial fluid)
- In the setting of miscarriage or stillbirth, isolation of *L. monocytogenes* from placental or fetal tissue

Case classification

Confirmed: a clinically compatible case that is laboratory-confirmed

Comment

The usefulness of other laboratory methods such as fluorescent antibody testing or polymerase chain reaction to diagnose invasive listeriosis has not been established.

Information Needed for Investigation

Verify the diagnosis. What laboratory tests were conducted and what were the results? **Establish the extent of illness**. Determine if household or other close contacts are, or have been ill, by contacting the health care provider, patient or family members. Determine if case is associated with food recall.

Contact the Regional Communicable Disease Coordinator, if an outbreak is suspected.

Case/Contact Follow Up And Control Measures

Determine the source of infection to prevent other cases:

Missouri Department of Health and Senior Services Communicable Disease Investigation Reference Manual



Division of Environmental Health and Communicable Disease Prevention								
Section: 4.0 Diseases and Conditions Updated 7/03								
Subsection: Listeriosis	Page 3 of 9							

- Does the person have contact with wild or domestic animals or work in animal operations?
- Have there been other cases linked by time, place or person?
- Has the person been exposed to a recalled food product?
- Has the person been exposed to nonpasteurized dairy products?

Control Measures

See the Listeriosis section of the <u>Control of Communicable Diseases Manual</u> (CCDM), "Control of patient, contacts and the immediate environment".

See Listeriosis section of the Red Book.

- Investigate outbreaks to identify a common source of infection, and prevent further exposure to that source.
- Pregnant women and immunocompromised individuals should avoid soft cheeses such as Brie, Camembert, and Mexican style cheeses. They should cook, until steaming hot, leftover foods or foods such as hot dogs. They should avoid deli meats and eat only properly cooked meats and pasteurized dairy products. They should also avoid contact with potentially infective materials, such as aborted animal fetuses on farms.
- Antimicrobial therapy of infection diagnosed during pregnancy may prevent fetal or perinatal infection and its consequences. (2)
- Ensure safety of foods of animal origin. Pasteurize all dairy products where possible. Irradiate soft cheeses after ripening or monitor nonpasteurized dairy products, such as soft cheeses, by culturing for *Listeria*.
- Processed foods that are found to be contaminated by *Listeria monocytogenes* (e.g., during routine bacteriologic surveillance) should be recalled.
- Thoroughly wash raw fruits and vegetables before eating.
- Thoroughly cook raw food from animal sources such as beef, pork, or poultry.
- Wash hands, knives, and cutting boards after handling uncooked foods.
- Avoid the use of untreated manure on food crops.

Laboratory Procedures

Human Specimens:

Diagnosis is based on culture of the organism from blood, cerebrospinal fluid or other normally sterile site. Initial testing is NOT provided by the State Public Health Laboratory (SPHL). However, the private laboratory can send an isolate of the cultured organism to the SPHL for confirmation. The SPHL does this testing for epidemiological purposes at no charge to the submitting laboratory.



Division of Environmental Health and Communicable Disease Prevention								
Section: 4.0 Diseases and Conditions Updated 7/03								
Subsection: Listeriosis	Page 4 of 9							

Food Samples:

Food samples can be sent to SPHL to be tested for *Listeria*. Samples should be collected in a sterile sample container and contain at least four ounces of the food to be tested. Please call the Regional Communicable Disease Coordinator prior to sending a sample.

Reporting Requirements

Listeriosis is a reportable disease and shall be reported to the local health authority or to the Missouri Department of Health and Senior Services (DHSS) within 3 days of first knowledge or suspicion by telephone, facsimile, or rapid communication.

- 1. For all reported cases, complete a "Disease Case Report" (CD-1), and a "Record of Investigation of Enteric Infection" (CD-2C) revised 6/02.
- 2. Entry of the completed CD-1 into the MOHSIS database negates the need for the paper CD-1 to be forwarded to the Regional Health Office.
- 3. Send the completed secondary investigation form to the Regional Health Office.
- 4. All outbreaks or "suspected" outbreaks must be reported as soon as possible (by phone, fax or e-mail) to the Regional Communicable Disease Coordinator. This can be accomplished by completing the Missouri Outbreak Surveillance Report (CD-51).
- 5. Within 90 days from the conclusion of an outbreak, submit the final outbreak report to the Regional Communicable Disease Coordinator.

References

- 1. Chin, James, ed. "Listeriosis." <u>Control of Communicable Diseases Manual</u>. 17th ed. Washington, DC: American Public Health Association, 2000: 296-299.
- American Academy of Pediatrics. "Listeria monocytogenes Infections (Listeriosis)." In: Pickering, LK., ed. 2000 Red Book: Report of the Committee on Infectious Diseases. 25th ed. Elk Grove Village, IL. 2000: 372-374.
- 3. Memorandum from the Centers for Disease Control and Prevention dated January 4, 2000. Nationally notifiable infectious disease list and case definition revisions: Update from the 1999 Council of State and Territorial Epidemiologists (CSTE) Meeting.

Other Sources of Information

- 1. Lorber, Bennett. "Listeria monocytogenes." Principles and Practice of Infectious Diseases. 5th ed. Eds. Gerald L. Mandell, John E. Bennett, and Raphael Dolin. New York: Churchill Livingstone, 2000: 2208-2215.
- 2. Armstrong, Donald and Gellin, Bruce G. "Listeria monocytogenes Infections." <u>Bacterial Infections of Humans Epidemiology and Control</u>. 3rd ed. Eds. Alfred S. Evans and Philip S. Brachman. New York: Plenum, 1998: 421-436.



Division of Environmental Health and Communicable Disease Prevention							
Section: 4.0 Diseases and Conditions Updated 7/03							
Subsection: Listeriosis	Page 5 of 9						

Web Sites

- 1. Centers for Disease Control and Prevention Listeriosis (also available in Spanish) http://www.cdc.gov/ncidod/dbmd/diseaseinfo/listeriosis_g.htm (28 July 2003)
- 2. Handbook on basic facts regarding foodborne pathogenic microorganisms and natural toxins http://vm.cfsan.fda.gov/~mow/intro.html (28 July 2003)
- 3. Karen B Weinstein. "*Listeria monocytogenes*." eMedicine Journal, Nov 19 2001, V 2, N 11 http://www.emedicine.com/MED/topic1312.htm (28 July 2003)

Listeriosis

Fact Sheet

What is listeriosis?

Listeriosis is a bacterial infection caused by *Listeria monocytogenes*. While many bacteria generally infect specific locations within the human body, listeria may infect many different sites, such as the brain or spinal cord membranes, or the bloodstream.

Who gets listeriosis?

Anyone can get the disease, but those at highest risk are newborns, the elderly, people with weakened immune systems and pregnant women. About 30 percent of cases occur in newborns within the first three weeks of life.

When do *listeria* infections occur?

Infections occur throughout the year. Although most cases occur sporadically, foodborne outbreaks do occur.

How is listeriosis spread?

Listeria bacteria are widely distributed in nature and can be found in water and soil. Infected animals may also serve as sources. Unlike other organisms, *listeria* can be spread by several different methods. Ingestion (foodborne transmission) of the organism, such as through unpasteurized milk or contaminated vegetables is often a source of cases. In newborn infections, the organism may be transmitted from mother to fetus in utero, or directly to the fetus at the time of birth. Direct contact with the organism can cause lesions on the hands or arms, and person-to-person transmission can occur through sexual contact. Infection is also possible by inhaling the organism.

What are the symptoms of listeriosis?

Because listeriosis can affect many different parts of the body, the symptoms vary. For meningoencephalitis, the onset can be sudden with fever, intense headache, nausea, vomiting and signs of meningeal irritation. In other body locations, various types of lesions at the site of infection are the primary symptom. In most cases, *listeria* infection causes fever and influenza-like symptoms resembling many other illnesses.

How soon after exposure do symptoms appear?

Listeriosis has an extremely variable incubation period. In large outbreaks, the range has extended from three to 70 days but usually within a month.

How is this disease diagnosed?

Specific laboratory tests are the only way to identify this disease. Since many cases may be mild, the disease may be more common than is realized.

Are there any unusual features of this disease?

Listeria infections are a significant risk for pregnant women, who may not experience obvious symptoms. Infection of the fetus can occur before delivery and can cause abortion as early as the second month of pregnancy, but more often in the fifth and six months. An infection later in pregnancy may cause exposure during birth, sometimes resulting in infection of the newborn, which may be fatal.

Does past infection with listeria make a person immune?

Past infection appears to produce some protective immunity.

What is the treatment for *listeria* infection?

Several antibiotics are effective against this organism. Ampicillin, either alone or in combination with other antibiotics, is frequently used.

What can be done to prevent the spread of this disease?

Since the organism is widespread in nature, basic sanitary measures such as using only pasteurized dairy products, eating cooked meats and washing hands thoroughly before preparing foods offer the best protection. Pregnant women and persons with weakened immune systems may wish to avoid such foods as soft cheeses and raw hot dogs. Although the risk of listeriosis associated with foods from deli counters is relatively low, pregnant women and immunosuppressed persons may choose to avoid these foods or thoroughly reheat cold cuts before eating.

Missouri Department of Health and Senior Services Section for Communicable Disease Prevention Phone: (866) 628-9891 or (573) 751-6113



MISSOURI DEPARTMENT OF HEALTH AND SENIOR SERVICES SECTION OF COMMUNICABLE DISEASE CONTROL AND VETERINARY PUBLIC HEALTH

RECORD OF INVESTIGATION OF ENTERIC ILLNESS

NAME: (LAST,	formation with st	naded title	s is not i	equired if	entered or		ATE OF		r entei	red into		RACE	
IANE: (LASI,	rikol, Wil)					'	/AIE UF	 	AGE:	GENL	JER:	RACE	
ARENT(S) NA	ME IF NOT ADULT:				PHONE NO.:		-						
HOME ADDRESS:					CITY:			STATE:		ZIP CODE: COUNTY:			
MPLOY	MENT / CHILD CA	RE (*See re	verse side	for High-Risk	Employment	info	rmatio	n.)					
LACE OF EMP		(ADDRESS:				,		PHON	NE NO.:		
OCCUPATION:					JOB DUTIES:								
OCCI ATION.					JOB BOTIES.								
CHOOL / CHIL	D CARE ATTENDED:				GRADE OR RO	OM:							
CHOOL / CHIL	D CARE ADDRESS:				CITY:				STATE	STATE: ZIP CODE:			
Symptom	IS:* (Check Yes or N	No and number	er the order	in which syn	nptoms first p	rese	nted)		•		•		
ORDER NO.	SYMPTOM	YES NO	ORDER NO.	SYM	PTOM	YES	NO	ORDER NO.		SYMPTOM		YES	_
	Nausea			Bloody Dia	ırrhea	<u> </u>	<u> </u>		Malaise			-	
	Vomiting Diarrhea			Cramps Chills					Heada Dizzine				<u> </u>
	Watery Diarrhea			Fever	0	Ħ			Other	,33			
Disease	,		I			_			1				_
IAGNOSIS:				ONSET DATE	/ TIME:*				DURATIO	ON OF SYMP			
IOUD ATION D	EDIOD.*			PHYSICIAN C	<u> </u>			am 🗌 pm	_	LUCODITAL	hr	S.	
ICUBATION P	ERIOD:				No No		DATE:	1 1		HOSPITAL Yes			
ROVIDER NAM	ME:		CITY:	1es				STATE		PHONE NO			
REATMENT: ((TYPE, AMOUNT)									DATE:*	,		
7.5		DATE OF DEA	TH:	CAUSE OF DEATH	1 :					,			
Recovere	ed Died	1 1	1										
Patient H	istory (Limit patient	responses to	within one	e disease inc	ubation period	l.)							
RAVEL: (OUT	SIDE OF HOME COMMUNITY	') DATE(S):*					LOCA	TION(S):					
OME WATER													
☐ Private (t					☐ Bottled W	/ater	(brand)					
	ater District (Name)				Other water								
IOME SEWAGI	E DISPOSAL SYSTEM:												
☐ Private (t	ype)				☐ Commun	ity Sy	/stem (Name)					
RECREATIONA	L WATER CONTACT: (SWIMI	MING POOL, LAKE	E, RIVER, ETC.)	Locati	on.							
☐ Yes ☐] No Type:												
ET / ANIMAL E	EXPOSURE: (DOMESTIC PET	TS, LIVESTOCK, C	THER)		24,55								
] Yes □		mals ill: 🔲 Ye			nimal Type(s):								
					ate(s)* of Anim	al Ex	posure):					
	imal Exposure: Animal Exposure:												
Comments:	Animai Exposure												
ood**		NAME			STREE	TADI	DESS			CI	TY / STATI	=	
<u> </u>		MAINE			SINE					- CI	, GIAII	=	
Grocery stor	es												
outinely use	ed:												
Restaurants													
outinely use													
					<u> </u>		<u> </u>						
	SOURCES: (e.g., ETHNIC, UN	IPASTEURIZED, H	OME CANNED)	TYPE / I	OCA.	ΓΙΟN:						

^{*} Epi Calendar (reverse side) may be used to help determine time periods.

** Attach separate 3-day food history if multiple cases are known/suspected.

Laboratory Tests*: Record Diagnostic Information in Section 41 of CD-1 Report and/or attach copy of lab slip(s)													
Are there other associated cases? ☐ Yes ☐ No					If yes, how	many?		How Associated:					
List ill contacts													
NAME & A	DOB / AGE	SEX		ATION TO TIENT	SIMILAR ILLNESS YES NO		ONSET DATE	CONF YES	LAB CONFIRMED		CD-1 AND ENTERIC FORM COMPLETED YES NO		
										NO			
High Risk Empl	oyment Inform	ation (e.g., Food	d Handl	er, Child C	are or Heal	th Care	Worker)					
DATE(S) WORKED PRIOR	TO ONSET OF ILLNESS	**					EXCLUDE ☐ Yes	D FROM WORK?		DATE:* /	1		
IF YES, BY WHOM:					TITLE:								
FOLLOW-UP SPECIMEN(S	B) REQUIRED? DAT	E COLLECTED:* /	1.	SULTS:*		2			3				
LAB:		WERE (DISCUSSED WI	ITH PATIE	NT?	BY:					
RETURNED TO WORK? ☐ Yes ☐ No	[DATE:* / /			EXPECTED (DATE:*	"		JDED FROM		SK DUTIES	5?	
SEXUAL PREFERENCE: Heterosexual RECREATIONAL DRUG US Yes No	☐ Homosexual SE: □	☐ Bisexual [DRUGS OF CHOICE:	_ Unkn	own [] N/A				PLE PARTI				
*Epi Calendar: MONTH(S) / DATES:		VEAD.			DIOTAGE:			L MODI	·-				
MONTH(S) / DATES:		YEAR:			DISEASE:			WORK	.:				
Sunday	Monday	Tuesday	-	Wedneso	lay	Thursday Friday _			Saturday				
					_								
Sunday	Monday	Tuesday	-	Wedneso	lay	Thurs	day	_ Friday		Satu	rday		
Sunday	Monday	Tuesday	-	Wedneso	lay	Thursday		_ Friday	_	Saturday		lay	
OTHER PERTINENT EPIDEMIOLOGICAL DATA (TO INCLUDE PROBABLE SOURCE):													
JILEN ENIMENT EPIDE	OLOGIOAL DATA (TO		JOJNOE)	•									
INVESTIGATOR:									DATE COM	PLETED:			
>													

MO 580-0802 (6-02)